## CANADIAN IMPORTATION OF CATTLE FROM EUROPE

Because of the possibility of introducing foot and mouth disease, movement of cattle from European countries to Canada has not been considered safe in past years. In the early 1960's, however, the improved disease situation in France and in Europe generally, together with technological advances in laboratory testing procedures, made possible the development of procedures to safeguard against the introduction of foot and mouth disease from countries not traditionally free of this disease.

In considering such an importation, Canada, as a country on a continent free of foot and mouth disease, recognized the need for effective disease control measures.

Cattle importations are governed by a permit system. The procedures established are rigid and involve inspection, testing and quarantine both in the country of origin and again on arrival in Canada. Cattle selected must not be over nine months of age on entering the Canadian quarantine station and must not have been vaccinated against foot and mouth disease. Cattle intended for export to Canada are inspected in the country of origin by a Canadian veterinary officer at assembly points, on the farms of origin and during the official quarantine. They are examined and certified free of diseases such as bluetongue, Johne's disease, leucosis, Vibrio fetus and Trichomonas infections. They are submitted to various tests to detect virus or antibodies of foot and mouth disease, as well as tests for brucellosis, tuberculosis, leucosis and leptospirosis. During the official European quarantine period of thirty days the cattle are again submitted to tests for foot and mouth and other diseases. All feed and litter used during this quarantine period, as well as during the sea voyage to Canada, is provided directly from Canada.

On arrival at the Canadian quarantine station, all tests are repeated and a test for bluetongue disease is also carried out. The Canadian quarantine station is at Grosse Ile, an island in the St. Lawrence river about thirty miles east of Quebec city. The station accommodates approximately 240 head of cattle and fully meets the requirements of a maximum security facility. During the Canadian quarantine, a period of at least ninety days, Canadian cattle, swine and sheep previously tested and found negative to the above diseases are continuously exposed to the quarantine animals as susceptible controls. The Canadian Government reserves the right to dispose of all or any imported or test animals that show clinical symptoms of disease, or which are not classed as negative to any of the tests.

Following their release from the Canadian quarantine station, the imported cattle undergo a further three-month quarantine period on the importer's farm in contact with his susceptible cattle.

A total of 565 cattle have been imported from France and Switzerland since the first importation in 1965.

Contributed by Health of Animals Branch, Canada Department of Agriculture.

# CANADIAN JOURNAL OF COMPARATIVE MEDICINE REVUE CANADIENNE DE MÉDECINE COMPARÉE

The Research Journal of the Canadian Veterinary Medical Association

La Revue de Recherche de l'Association Canadienne des Médecins Vétérinaires

## EDITORIAL BOARD

**Editor-in-Chief:** J. F. Frank, Hull, Que. — **Managing Editor:** Miss Nella McKellar, Ottawa, Ontario — **Associate Editors:** R. Dunlop, Saskatoon, Saskatchewan; Joan Budd, Guelph, Ontario; J. G. Lafortune, St. Hyacinthe, Quebec. — **Editor Emeritus:** C. A. Mitchell, Ottawa, Ontario.

### INSTRUCTIONS FOR AUTHORS

Research articles in English or French pertaining to comparative medicine will be considered, provided they are not being submitted elsewhere. All manuscripts will be subjected to editorial action and to review for adequacy of scientific content.

Manuscripts should be in duplicate typewritten (double spaced), with 1 inch margins. An abstract should be included. References should be listed alphabetically and referred to in the text by arabic numerals in brackets.

A publication charge of \$20. per printed page, plus cost of engravings, is payable by contributors. Instructions for ordering reprints will be sent with the galley proofs.

Published quarterly by the Canadian Veterinary Medical Association.

Correspondence should be addressed to: —

Canadian Journal of Comparative Medicine, 360 Bronson Ave., Ottawa 4, Ontario.

### Subscription rates:

### INSTRUCTION AUX AUTEURS

La Revue publie des articles originaux en français ou en anglais, traitant de médecine comparée. Le comité de rédaction procède à une évaluation du contenu scientifique des manuscrits qui lui sont soumis et suggère les modifications requises pour l'édition. L'original et une copie du manuscrit doivent être soumis, dactylographiés à double interligne, avec une marge d'un pouce à gauche de la page. Un résumé doit accompagner chaque article. Les références bibliographiques seront classifiées par ordre alphabétiques à la fin de l'article, et indiquées dans le texte au moyen de chiffres arabes entre parenthèses.

Les auteurs doivent assumer les frais de publication; le tarif est de \$20.00 par page imprimée, en plus du coût des figures et des planches. Un formulaire de commande de tirés-à-part sera adressé avec les épreuves.

Publication trimestrielle de l'Association canadienne des vétérinaires.

Adresser toute correspondance à:—
Revue canadienne de Médecine
comparée,
360 Ave. Bronson,
Ottawa 4, Ontario.

Authorized as second class mail Post Office Department, Ottawa, and for payment of postage in cash.

Formerly published by National Business Publication Ltd., Gardenvale, Quebec.

Vol. 32 — July, 1968

PROPERTY OF US ARMY
TECHNING FORWARD ARSENAL, MD.

DIFFERENTIATION OF DUCK PLAGUE, NEWCASTLE DISEASE, FOWL PLAGUE, AND **DUCK HEPATITIS VIRUSES** 

Serums from 5 duck-plague-convalescent ducks and antiserums produced against Newcastle disease, fowl plague, and duck hepatitis were tested for their ability to inhibit the formation of plaques by HDPAV. The cultures were inoculated with virusserum mixtures constituted to contain approximately 55 PFU of HDPAV in a 0.2ml and final serum dilution varying in 2fold increments from 1:2 to 1:16 (Table III). Plaques were inhibited in the culture only when the homologous serum was employed. The plaques were equal in number to those plaques developed in the cultures that were inoculated with mixtures of virus and normal duck serum. These tests indicated that the plaque technique was applicable for the identification of duck plaque virus and its differentiation and nonrelationship to Newcastle disease, fowl plague, and duck hepatitis viruses.

### DISCUSSION

The plaque technique proved most useful when it was applied to viruses passed either in ducks, chicken- or duck-embryocell culture. The plaques were indistinct when duck-liver tissue suspensions were employed. However, when liver tissue was passed in DE, the efficiency of plaque development increased. Using a suspension of chorio-allantoic membranes infected with the liver suspension, distinct plaques developed, but they were smaller in size than those obtained with virus from the 5th cell culture passage. The plaques obtained from this passage were small and medium in size. Those produced from the attenuated virus were uniform and of medium size. Morphologically, they were indistinguishable from those produced by the cell-passed virulent virus.

The plaque technique was used to assay the viruses and to determine their antigenic relationships. It was established by experiments concerned with neutralization of the viruses, using the plaque inhibition technique, that the DPLIVV and HDPAV were serologically identical and that they were not related to Newcastle disease, fowl plague, and duck hepatitis viruses.

### **ACKNOWLEDGEMENTS**

The assistance rendered by Dr. R. Trautman in the computation of portions of the data collected and the excellent technical assistance of Mrs. Joan Grohoski are gratefully acknowledged.

### REFERENCES

- DARDIRI, A. H., W. R. HESS, S. S. BREESE JR., and H. R. SEIBOLD. Characterization of duck plague virus from a duck disease outbreak in the United States. Presented at the 39th Northeastern Conference on Avian Diseases, State Univ. of New York, June 19-21, 1967, Stony Brook, New York.
   DULBECCO, R., and M. VOGT. Plaque formation and isolation of pure lines with poliomyelitis virus. J. exp. Med. 99: 167-182. 1954.
   JANSEN, J. Duck plague. Br. vet. J. 117: 349-356. 1961.

- 1961.
  4. SCHMIDT, NATHALIE J. Tissue culture methods and procedures for diagnostic virology. In "Diagnostic Procedures for Virus and Rickettsial Diseases".
  3rd ed. New York: American Public Health Association, Inc., 1964. pp. 101-102.

### NOTICE

New address for the C.J.C.M. Office:

360 Bronson Ave.. Ottawa 4, Canada.

### AVIS

Nouvelle adresse du bureau de la Revue Canadienne de Médecine Comparée:

> 360 avenue Bronson, Ottawa 4, Canada.

### **ADVERTISERS' INDEX**

Canada Department of Agriculture IFC
Hill Packing Company VI
Merck, Sharp and Dohme of Canada Limited IV
Pitman-Moore Limited IBC
Wild of Canada Ltd OBC
Rougier Incorporated VI

Address correspondence re advertising to: Canadian Journal of Comparative Medicine, 360 Bronson Ave., Ottawa 4, Canada.

### **GUIDELINES FOR CONTRIBUTORS AND THEIR SECRETARIES**

In order to facilitate the submission of manuscripts in a form which permits rapid and efficient editing, and to ensure uniformity of style, the C.J.C.M. has drawn up the following outline. It is felt that reference to these guidelines, and to a recent issue of the Journal itself, will enable secretaries to draft manuscripts requiring a minimum of structural change upon receipt.

### **GENERAL**

Manuscripts written in either English or French will be considered.

All parts of the manuscript, including footnotes, tables and illustrations, should be submitted on white paper, not flimsy, of a size  $8\frac{1}{2}$  x 11 inches.

All typing should be at least double-spaced and with a minimum of one-inch margins.

The original and one clear duplicate must be submitted.

Do not underline unless the type is to be set in italics.

Consult a recent issue for the format of headings, sub-headings, footnotes, etc.

Standard abbreviations only such as mg, ml, hr, etc. are acceptable without definition. Less common abbreviations should be defined by writing the words out in full when first used.

Words, spelling and usage should conform to Webster's Collegiate Dictionary or Le Dictionnaire Encyclopédique Quillet (6 tomes).

Numerals expressing measurement or quantities must be written using decimals, as 0.1 gm, 1.0 ml, 10.0 mm, etc. Numbers from one to ten should be written out. Larger numbers are to be expressed in Arabic numerals.

### REFERENCES

Type references double-spaced.

Use only abbreviations sanctioned by the "World List of Scientific Periodicals".

Identify references in the text by Arabic numerals.

Check the text to see that all references in the list are referred to.

List all references alphabetically by surname of the first author listed.

The acceptable style, with respect to sequence and punctuation marks, must be followed. If in doubt, consult a recent issue, but the following example will summarize the desirable style:

14. LEWIS, V., and J. B. BROOKS.

Comparison of fluorochromes for
the preparation of fluorescent-anti-

body reagents. J. Bact. 88: 1520-1521, 1964.

STABLEFORTH, A. W. and I. A. GALLOWAY. Infectious diseases of animals. Volume 2. London: Butterworths Scientific Publications. 1959.

### **TABLES**

Consult a recent issue to see how tables are designed and how lines are placed.

Do not include vertical lines.

Number tables with Roman numerals.

Designate footnotes by symbols.

Type tables on separate pages and always type double-spaced.

Place tables together in the manuscript after the text and continue the page numbers

### **FOOTNOTES**

Designate footnotes by symbols. Insert footnotes at the bottom of the proper page, below a solid horizontal line. Type footnotes for tables below the appropriate table.

### **ILLUSTRATIONS**

All illustrations and figures should be numbered consecutively in Arabic numerals, and should be similarly referred to in the text.

All should bear a title.

Photographs should be unmounted glossy prints with sharp contrast. They should be trimmed so that essential features only are shown.

Each illustration must be clearly marked on the back with the appropriate figure number, name of author, and title of the article.

Drawings and charts should be in black India ink on white paper.

Letters, numerals and all necessary lines must also be in ink, not typewritten.

The originals and one set of clear copies (e.g. small photographs) are to be submitted.

A scale must be submitted with all photomicrographs and with other photographs where reference to size is made.